

Next-to-next-to-leading order corrections to deeply virtual Compton scattering: the non-singlet case

D. Müller

*Institut für Theoretische Physik, Universität Regensburg
D-93040 Regensburg, Germany*

*Department of Physics and Astronomy, Arizona State University
P.O. Box 871504, Tempe, AZ 85287-1504, USA*

Abstract

We study perturbative QCD corrections to deeply virtual Compton scattering on an unpolarized nucleon target in the flavor non-singlet sector to next-to-next-to-leading order accuracy, restricting ourselves to the kinematically dominant amplitude. The difference between the standard minimal subtraction and the conformal scheme, in which conformal symmetry is manifest, is studied to next-to-leading order. Beyond this order we employ conformal symmetry for the evaluation of perturbative corrections. Within a certain class of generalized parton distributions we find moderate radiative corrections.

Keywords: perturbative corrections, deeply virtual Compton scattering, conformal symmetry

PACS numbers: 11.10.Hi, 12.38.Bx, 13.60.Fz

1 Introduction

The hard-exclusive photo electroproduction off a nucleon, among a class of hard exclusive but inelastic processes, is generally considered as the theoretic cleanest process to gain access to the so-called generalized parton distributions (GPDs) [1, 2]. These non-perturbative distributions contain manifold information that cannot be gained from measurements of deep (semi-)inelastic or elastic and other exclusive processes [3]. Based on a partonic interpretation, GPDs allow us to reveal the internal structure of the probed hadron, especially of the nucleon, from a new perspective, for comprehensive reviews see Ref. [4].

We recall that the process in question has two interfering subprocesses, namely, the Bethe-Heitler bremsstrahlung and deeply virtual Compton scattering (DVCS). The former is known in terms of the electromagnetic form factors and so the latter can be measured in several asymmetries, see e.g., Ref. [5], that appear in leading order of the expansion with respect to the inverse photon virtuality, i.e., single and double spin and charge asymmetries [6, 7, 8, 9]. Relying on the validity of the operator product expansion (OPE) of two electromagnetic currents, the DVCS amplitude factorizes to leading power, i.e., twist-two, in short- and long-distance physics, where the former and latter are incorporated in (resummed) Wilson-coefficients and GPDs, respectively [1]. Equivalently, in the partonic framework, the DVCS amplitude to leading power accuracy is represented as convolution of hard-scattering amplitudes, systematically calculable in powers of the strong coupling, and GPDs. The factorization of collinear singularities was shown to all orders in perturbation theory [10].

Let us remind that for experimental accessible scales, the perturbative description of elastic exclusive processes is controversially debated [11]. With respect to this fact one might wonder whether the perturbative GPD framework is justified. For the GPD phenomenology it is essential to clarify this issue; unfortunately, this is an interlaced task. The relative size of perturbative and non-perturbative, i.e., power suppressed contributions, might serve as criteria for the reliability of the perturbative framework. The DVCS hard-scattering amplitudes (Wilson-coefficients) were perturbatively evaluated up to next-to-leading order (NLO) in the strong coupling [12, 13]. It turned out that these radiative corrections can be of the order of 30% to 50% for fixed target kinematics [14]. Even much larger corrections have been reported for the kinematics of collider experiments [15]. Mainly, but not only, this is due the appearances of gluonic GPDs at NLO, see also discussion in Sect. 6.2.3 of Ref. [8]. The problem arises whether these sizeable corrections are related to the appearance of gluons, are induced by an awkward choice of GPD ansätze, or indicate that at experimental accessible scales the perturbative regime has not be fully set in.

To get a deeper insight into this issue, we study in this letter perturbative corrections to the DVCS process beyond NLO accuracy within a framework that avoids a cumbersome diagrammatic

calculation. Based on the conformally covariant OPE (COPE), the perturbative corrections up to next-to-next-to-leading order (NNLO) can be economically evaluated [16, 12], see also review [17]. Thereby we restrict ourselves to the flavor non-singlet part of the helicity conserved twist-two amplitude in the parity even sector, which is the dominant contribution in several DVCS observables. In Sect. 2 we outline the evaluation of this amplitude by means of the COPE [16, 12, 18]. This yields a divergent series of conformal GPD moments, which is resummed within a Mellin-Barnes integral. In Sect. 3 we numerically study the radiative corrections, mainly due to the Wilson-coefficients, at NLO and NNLO. Especially, we explore the numerical differences of the NLO corrections in the modified minimal subtraction ($\overline{\text{MS}}$) and conformal subtraction (CS) schemes. We then provide estimates to NNLO accuracy. Finally, in Sect. 4 we conclude.

2 General formalism

Let us first recall the standard perturbative QCD framework. Usually, one employs the $\overline{\text{MS}}$ scheme, which is based on dimensional regularization and the removal of the poles with respect to the dimensional parameter $(4 - n)/2$. This scheme is used twofold: (i) to define the renormalized strong coupling and (ii) for the factorization of collinear singularities or if one wishes for the renormalization of (leading twist) composite operators¹, which are labelled by the good quantum number spin. In exclusive reactions such operators with given spin are also be build within total derivatives. In general operators with the same spin will mix under renormalization. To get rid of this mixing phenomenon, one changes at some stage of the full calculation in an explicit or implicit manner to a basis of multiplicatively renormalizable operators, i.e., one diagonalizes the evolution equation [19]. For vanishing $\beta(\alpha_s)$ function this transformation leads to conformal operators [20]. Such operators transform covariantly under the so-called collinear conformal transformation $SL(2, \mathbb{R})$ and are members of infinite dimensional conformal multiplets (towers) that are labelled by the conformal spin. With other words one takes explicitly or implicitly advantages from the underlying conformal symmetry of the classical QCD Lagrangian. If the trace anomaly of the energy-momentum tensor, proportional to $\beta(\alpha_s)$, is absent, conformal symmetry is present in perturbative QCD and its predictive power can be employed at any order.

In the following we evaluate the scattering amplitude for the DVCS process, which is expressed in terms of the time ordered product of two electromagnetic currents

$$T_{\mu\nu}(q, P_1, P_2) = \frac{i}{e^2} \int dx e^{ix \cdot q} \langle P_2, S_2 | T j_\mu(x/2) j_\nu(-x/2) | P_1, S_1 \rangle, \quad (1)$$

¹To leading twist approximation there is an one-to-one cross talk between the ultra-violet and collinear singularities in the partonic matrix elements of these operators.

where $q = (q_1 + q_2)/2$ (μ and q_2 refers to the outgoing real photon). The incoming photon has a large virtuality $q_1^2 = -Q^2$ and requiring that in the limit $-q^2 = Q^2 \rightarrow \infty$ the scaling variables

$$\xi = \frac{Q^2}{P \cdot q}, \quad \eta = -\frac{\Delta \cdot q}{P \cdot q}, \quad P = P_1 + P_2, \quad \Delta = P_2 - P_1. \quad (2)$$

and the momentum transfer squared Δ^2 are fixed, the dominant contributions arise from the light-cone singularities of the time ordered product. In this generalized Bjorken limit one can now employ the OPE to evaluate the hadronic tensor in terms of the leading twist-two operators, where for DVCS kinematics $\eta \simeq \xi$ and $Q^2 \simeq Q^2/2$.

Before we outline this step let us introduce a parameterization of the hadronic tensor

$$T_{\mu\nu}(q, P, \Delta) = -\tilde{g}_{\mu\nu} \frac{q_\sigma V^\sigma}{P \cdot q} - i\tilde{\epsilon}_{\mu\nu\rho\sigma} \frac{q^\rho A^\sigma}{P \cdot q} + \dots \quad (3)$$

To ensure current conservation, the metric and Levi-Civita tensors are contracted here with projection operators, for explicit definitions of $\tilde{g}_{\mu\nu}$ and $\tilde{\epsilon}_{\mu\nu\rho\sigma}$ see, e.g., Ref. [7]. The ellipsis indicates terms that are finally power suppressed in the DVCS amplitude or a determined by the gluon transversity GPD, which is not considered here. We note that in the forward limit $\Delta \rightarrow 0$ the first and second term on the r.h.s. are expressed by the deep inelastic structure functions F_1 and g_1 . In the parity even sector the vector

$$V^\sigma = \bar{U}(P_2, S_2) \left(\mathcal{H} \gamma^\sigma + \mathcal{E} \frac{i\sigma^{\sigma\rho} \Delta_\rho}{2M} \right) U(P_1, S_1) + \dots, \quad (4)$$

is decomposed in Compton form factors (CFFs) $\mathcal{H}$ and $\mathcal{E}$, similar for the axial-vector A^σ in terms of $\tilde{\mathcal{H}}$ and $\tilde{\mathcal{E}}$, where again higher twist contributions are neglected.

Now we are in the position to employ the OPE. Let us first suppose that the trace anomaly is absent and so conformal symmetry can be employed. Formally, this can be achieved by assuming that a hypothetical fixed point exist, i.e., $\beta(\alpha_s^*) = 0$. We can then use the COPE, which tells us how the total derivatives are arranged. Moreover, it can be considered as a partial wave expansion with respect to the conformal spin $j + 2$. For the time-ordered product of two electromagnetic currents it reads in the flavor non-singlet sector as [21, 16]

$$\begin{aligned} \{T j_\mu(x) j_\nu(0)\}^{\text{NS}} &= \frac{1}{\pi^2} \frac{-\tilde{g}_{\mu\nu}}{(-x^2 + i0)^2} \sum_{j=1}^{\infty} [1 - (-1)^j] \frac{\Gamma(2 - \gamma_j/2)(2 + j + \gamma_j/2)}{2^{\gamma_j/2} \Gamma(j + 1 + \gamma_j/2)} C_j^{\text{NS}}([-x^2 + i0]\mu^2) \\ &\quad \times (-2ix_-)^{j+1} \int_0^1 du [u(1-u)]^{j+1+\gamma_j/2} \mathcal{O}_j^{\text{NS}}(ux_-) + \dots, \end{aligned} \quad (5)$$

where $x_- = \tilde{n} \cdot x$ with $\tilde{n}^2 = 0$ is the projection of the vector x on the light cone. The anomalous dimensions of the multiplicatively renormalizable operators $\mathcal{O}_j^{\text{NS}}$ are denoted as $\gamma_j(\alpha_s^*)$. The advantage of the COPE is that the Wilson-coefficients (electrical charge factors will be omitted)

$$C_j^{\text{NS}}(-x^2\mu^2) = (-\mu^2 x^2)^{\gamma_j/2} \frac{2^{j+1+\gamma_j/2} \Gamma(5/2 + j + \gamma_j/2)}{\Gamma(3/2) \Gamma(3 + j + \gamma_j/2)} c_j(\alpha_s^*), \quad (6)$$

are up to the normalization $c_j(\alpha_s^*)$ known. The conformal operators

$$\mathcal{O}_j^{\text{NS}}(ux_-) = \frac{\Gamma(3/2)\Gamma(1+j)}{2^j\Gamma(3/2+j)} (i\partial_+)^j \bar{\psi}(ux_-)\gamma_+ C_j^{3/2}\left(\frac{\vec{D}_+}{\partial_+}\right) \psi(ux_-), \quad (7)$$

are defined with a non-standard normalization. Here $\vec{D}_+ = \vec{D}_+ - \overleftarrow{D}_+$ and $\partial_+ = \vec{D}_+ + \overleftarrow{D}_+$ are the covariant and total derivatives, contracted with the light-like vector n , i.e., $n^2 = 0$ and $n \cdot \tilde{n} = 1$, $C_j^{3/2}$ are the Gegenbauer polynomials of order j with index $3/2$. The normalization is chosen so that in the forward limit $\Delta \rightarrow 0$ the reduced matrix elements of the conformal operators

$$\frac{1}{P_+^{j+1}} \langle P_2, S_2 | \mathcal{O}_j(0) | P_1, S_1 \rangle = \frac{1}{P_+} \bar{U}(P_2, S_2) \left(H_j(\eta, \Delta^2, \mu^2) \gamma_+ + E_j(\eta, \Delta^2, \mu^2) \frac{i\sigma_{+\nu} \Delta^\nu}{2M} \right) U(P_1, S_1) \quad (8)$$

coincide with the Mellin moments of parton densities. Especially, for odd values of j the forward limit of H_j is the sum of the quark and anti-quark density moments, i.e., $\lim_{\Delta \rightarrow 0} H_j = q_j + \bar{q}_j$. Moreover, in this limit the u -integration in Eq. (5) can be performed and leads to the well-known OPE that is used for the deep inelastic scattering structure function F_1 . Hence, $c_j(\alpha_s^*)$ are identified as the Wilson-coefficients of F_1 , known to NNLO order.

Employing the COPE (5), we can now achieve the factorization of the CFFs (4) in a straight-forward manner. Plugging Eq. (4) into the definition of the hadronic tensor (1), performing Fourier transform, and form factor decomposition (4), we arrive for the DVCS kinematics $\eta = \xi$ at [16]

$$\left\{ \begin{array}{c} \mathcal{H}^{\text{NS}} \\ \mathcal{E}^{\text{NS}} \end{array} \right\} = \sum_{j=1}^{\infty} \frac{1 - (-1)^j}{2} \xi^{-j-1} C_j^{\text{NS}}(\mu^2/Q^2) \left\{ \begin{array}{c} H_j^{\text{NS}} \\ E_j^{\text{NS}} \end{array} \right\} (\eta, \Delta^2, \mu^2) \Big|_{\eta=\xi}. \quad (9)$$

Here H_j^{NS} and E_j^{NS} , cf. Eq. (8), are given by the conformal moments of the corresponding GPDs

$$\left\{ \begin{array}{c} H_j^{\text{NS}} \\ E_j^{\text{NS}} \end{array} \right\} (\eta, \Delta^2, \mu^2) = \frac{\Gamma(3/2)\Gamma(1+j)}{2^j\Gamma(3/2+j)} \int_{-1}^1 dx \eta^j C_j^{3/2}\left(\frac{x}{\eta}\right) \left\{ \begin{array}{c} H^{\text{NS}} \\ E^{\text{NS}} \end{array} \right\} (x, \eta, \Delta^2, \mu^2). \quad (10)$$

Note that these expectation values are measurable on the lattice [22].

Unfortunately, the series (9) is divergent for $|\xi| < 1$ and must be resummed, e.g., by means of the Sommerfeld-Watson transformation [23]. Alternatively, using the trick that η is not equated to ξ allows to deal with Eq. (9) in the analogous manner as it is known from deep inelastic scattering. Namely, a dispersion relation allows to express the Mellin moments of the imaginary part by the partial waves, appearing in the COPE (9), and thus the inverse Mellin transform provides the imaginary part of the CFFs. The real part can be restored from the dispersion relation, too, and finally equating $\eta = \xi$ leads to the same representation as presented in Ref. [23]:

$$\left\{ \begin{array}{c} \mathcal{H}^{\text{NS}} \\ \mathcal{E}^{\text{NS}} \end{array} \right\} (\xi, \Delta^2, \mu^2) = \frac{1}{2i} \int_{c-i\infty}^{c+i\infty} dj \xi^{-j-1} \left[i + \tan\left(\frac{\pi j}{2}\right) \right] C_j^{\text{NS}}(Q/\mu) \left\{ \begin{array}{c} H_j^{\text{NS}} \\ E_j^{\text{NS}} \end{array} \right\} (\xi, \Delta^2, \mu^2), \quad (11)$$

where all singularities of the conformal GPD moments and conformal moments lie on the l.h.s. of the integration path ($c < 1$). Note, however, that the analytic continuation of the Wilson-coefficients (6) with respect to the conformal spin, analogously done as in deep inelastic scattering, leads essentially to an exponential 2^j growing with increasing j . For $\xi > 1$, this must be weighed down by a suppression factor that comes from the conformal moments (10). This is a rather nontrivial requirement on their analytic continuation. It should be done in such a way that the integral (11) remains unchanged if the integration contour is closed by an infinite arc, surrounding the first and forth quadrant. The residue theorem states then that this Mellin–Barnes integral is for $\xi > 1$ equivalent to the series (9). We note that the form of the integrand (11) does in fact not rely on conformal symmetry and so it can be used in any scheme within the corresponding Wilson-coefficients.

In the $\overline{\text{MS}}$ scheme the conformal symmetry is not manifestly implemented in the OPE, however, this failure can be cured by a finite renormalization [16, 12]. As explained above, in such a CS scheme and for $\beta = 0$ we can simply borrow the Wilson-coefficients $c_j(\alpha_s)$ and anomalous dimensions $\gamma_j(\alpha_s)$ from deep inelastic scattering. The inclusion of β proportional terms in Eq. (11) is, as factorization by itself, conventional. Two possibilities have been discussed in Ref. [18]. Namely, in the CS scheme, one might add the β proportional term from the $\overline{\text{MS}}$ scheme, evaluated to NNLO in Ref. [24], while in the $\overline{\text{CS}}$ scheme the running of the coupling is implemented in the form of the COPE (5) in such a way that the Wilson-coefficients autonomously evolve in the considered order [18]. In this letter we prefer the latter convention and, moreover, will equate the factorization and renormalization scales, i.e., $\mu = \mu_f = \mu_r$. After expansion with respect to α_s , we write, discarding in the following the superscript NS, the Wilson-coefficients (6) as

$$C_j = \frac{2^{j+1}\Gamma(j+5/2)}{\Gamma(3/2)\Gamma(j+3)} \left[C_j^{(0)} + \frac{\alpha_s(\mu)}{2\pi} C_j^{(1)}(\mathcal{Q}/\mu) + \frac{\alpha_s^2(\mu)}{(2\pi)^2} C_j^{(2)}(\mathcal{Q}/\mu) + \mathcal{O}(\alpha_s^3) \right], \quad (12)$$

$$C_j^{(0)} = 1, \quad C_j^{(1)} = c_j^{(1)} + \frac{\gamma_j^{(0)}}{2} \left[s_j^{(1)} + \ln \frac{\mu^2}{Q^2} \right], \quad (13)$$

$$C_j^{(2)} = c_j^{(2)} + \frac{\gamma_j^{(1)}}{2} \left[s_j^{(1)} + \ln \frac{\mu^2}{Q^2} \right] + \frac{\gamma_j^{(0)}}{2} \left[c_j^{(1)} \left\{ s_j^{(1)} + \ln \frac{\mu^2}{Q^2} \right\} + \frac{\gamma_j^{(0)}}{4} \right. \\ \left. \times \left\{ s_j^{(2)} + 2s_j^{(1)} \ln \frac{\mu^2}{Q^2} + \ln^2 \frac{\mu^2}{Q^2} \right\} \right] - \frac{\beta_0}{2} \left[c_j^{(1)} + \frac{\gamma_j^{(0)}}{2} s_j^{(1)} + \frac{\gamma_j^{(0)}}{4} \ln \frac{\mu^2}{Q^2} \right] \ln \frac{\mu^2}{Q^2}, \quad (14)$$

where

$$\gamma_j^{(0)} = C_F \left(4S_1(j+1) - 3 - \frac{2}{(j+1)(j+2)} \right), \quad C_F = \frac{4}{3}, \quad (15)$$

$$c_j^{(1)} = C_F \left[S_1^2(1+j) + \frac{3}{2}S_1(j+2) - \frac{9}{2} + \frac{5-2S_1(j)}{2(j+1)(j+2)} - S_2(j+1) \right], \quad (16)$$

$$s_j^{(1)} = S_1(j+3/2) - S_1(j+2) + 2\ln(2), \quad s_j^{(2)} = \left(s_j^{(1)} \right)^2 - S_2(j+3/2) + S_2(j+2). \quad (17)$$

The analytic continuation of the harmonic sums are defined by $S_1(z) = d \ln \Gamma(z+1)/dz + \gamma_E$ and $S_2(z) = -d^2 \ln \Gamma(z+1)/dz^2 + \zeta(2)$, where γ_E is the Euler constant. The first expansion coefficient of $\beta(g)/g = (\alpha_s/4\pi)\beta_0 + \mathcal{O}(\alpha_s^2)$ is $\beta_0 = (2/3)n_f - 11$, where n_f is the number of active quarks. The two-loop quantities $c_j^{(2)}$ and $\gamma_j^{(1)}$ are lengthy and can be obtained from Ref. [25, 26]. The evolution of the flavor non-singlet (integer) conformal moments in this $\overline{\text{CS}}$ scheme is governed by

$$\mu \frac{d}{d\mu} \left\{ \frac{H_j^{\text{NS}}}{E_j^{\text{NS}}} \right\} (\eta, \Delta^2, \mu^2) = - \left[\frac{\alpha_s(\mu)}{2\pi} \gamma_j^{(0)} + \frac{\alpha_s^2(\mu)}{(2\pi)^2} \gamma_j^{(1)} + \frac{\alpha_s^3(\mu)}{(2\pi)^3} \gamma_j^{(2)} + \mathcal{O}(\alpha_s^4) \right] \left\{ \frac{H_j^{\text{NS}}}{E_j^{\text{NS}}} \right\} (\eta, \Delta^2, \mu^2) - \frac{\beta_0}{2} \frac{\alpha_s^3(\mu)}{(2\pi)^3} \sum_{k=0}^{j-2} \eta^{j-k} \left[\Delta_{jk}^{\overline{\text{CS}}} + \mathcal{O}(\alpha_s) \right] \left\{ \frac{H_k^{\text{NS}}}{E_k^{\text{NS}}} \right\} (\eta, \Delta^2, \mu^2), \quad (18)$$

where the mixing matrix $\Delta_{jk}^{\overline{\text{CS}}}$ is not completely known.

We remark that the NLO corrections in the $\overline{\text{MS}}$ scheme can be easily evaluated from the conformal moments of the hard-scattering amplitude, e.g., given in Ref. [14]. All integrals, which are needed, are given in Appendix C of Ref. [18] for integer conformal spin. The analytic continuation is straightforward and so in the $\overline{\text{MS}}$ scheme Eq. (13) is to replace by

$$c_j^{\overline{\text{MS}}(1)} = C_F \left[2S_1^2(1+j) - \frac{9}{2} + \frac{5 - 4S_1(j+1)}{2(j+1)(j+2)} + \frac{1}{(j+1)^2(j+2)^2} \right] + \frac{\gamma_j^{(0)}}{2} \ln \frac{\mu^2}{Q^2}. \quad (19)$$

In this scheme also the complete anomalous dimension matrix is known to two-loop accuracy [20].

3 Numerical results

In this Section we numerically study the radiative corrections to the CFF

$$\mathcal{H}^{\text{N}^P\text{LO}} = \sum_{p=0}^P \left(\frac{\alpha_s(\mu)}{2\pi} \right)^p \frac{1}{2i} \int_{c-i\infty}^{c+i\infty} dj \xi^{-j-1} \frac{2^{j+1} \Gamma(5/2+j)}{\Gamma(3/2) \Gamma(3+j)} \left[i + \tan\left(\frac{\pi j}{2}\right) \right] c_j^{(p)} \left(\frac{Q}{\mu} \right) H_j^{\text{NS}}(\xi, \Delta^2, \mu^2) \quad (20)$$

at NLO ($P = 1$) and NNLO ($P = 2$) accuracy. Certainly, the size of the radiative corrections depend on the GPD distribution itself. As a GPD model assumption, we suppose that the expansion of H_j^{NS} in powers of ξ^2 induces a systematic expansion of the CFF (20). A closer look to this issue has been given in Sect. 4 of Ref. [23]. The GPD moments are generically parameterized as

$$H_j^{\text{NS}}(\Delta^2, \xi, \mu_0^2) = F^{\text{NS}}(\Delta^2) \frac{B(1 - \alpha(\Delta^2) + j, \beta + 1)}{B(1 - \alpha(\Delta^2), \beta + 1)} + \mathcal{O}(\xi^2). \quad (21)$$

In the forward limit, i.e., $\Delta \rightarrow 0$, these moments reduce to the Mellin moments of the unpolarized parton density. Hence, the parameters α and β characterize the small and large x behavior, respectively, i.e., $q(x, \mu_0) \propto x^{-\alpha}(1-x)^\beta$. For non-singlet parton densities the generic values are




Figure 1: The relative radiative corrections, defined in Eq. (22), are plotted versus ξ for the modulus [(a) and (b)] and phase [(c) and (d)] of the CFF (20) with $\alpha = 0.5$ [(a) and (c)] and $\alpha = -0.1$ [(b) and (d)]: NNLO in full (solid) and $\beta_0 = 0$ (dash-dotted) as well as in NLO for the $\overline{\text{CS}}$ (dashed) and $\overline{\text{MS}}$ (dotted) scheme. We always set $\mu = Q$ and $\alpha_s(Q^2 = 2.5 \text{ GeV}^2)/\pi = 0.1$.

$\alpha(\Delta^2 = 0) = 1/2$ and $\beta = 3$. In the off-forward kinematics we consider $\alpha(\Delta^2) = \alpha(0) + \alpha'(0)\Delta^2$ as a linear meson Regge trajectory with $\alpha(0) = 1/2$ and $\alpha'(0) = 0.9 \text{ GeV}^{-2}$. Note also that the conformal GPD moment with $j = 0$ reduces to the elastic form factor $F^{\text{NS}}(\Delta^2)$. Certainly, a more realistic ansatz of GPD moments is formed by a linear combination of the building blocks (21).

Let us first compare the radiative corrections in the $\overline{\text{MS}}$ and $\overline{\text{CS}}$ scheme to NLO accuracy. Strictly spoken, at a given input scale Q_0 there is no difference between both predictions, if the non-perturbative quantities are transformed, too, and a consequent expansion in α_s is performed. Usually, the GPD moments are taken from some non-perturbative (model) calculation or ansatz, where the matching with the perturbative prescriptions has its own uncertainties. So let us take the same input in both schemes and study the relative changes

$$K_{\text{abs}}^P = \frac{|\mathcal{H}^{\text{N}^P\text{LO}}|}{|\mathcal{H}^{\text{N}^{P-1}\text{LO}}|}, \quad K_{\text{arg}}^P = \frac{\arg(\mathcal{H}^{\text{N}^P\text{LO}})}{\arg(\mathcal{H}^{\text{N}^{P-1}\text{LO}})} \quad (22)$$

to the modulus and phase of the CFF (20). These quantities do not suffer from large radiative corrections as it is artificially the case for the real part of the amplitude². One should bear in

²The real part possesses a zero in the valence quark region, which position depends on the approximation. Hence, in the vicinity of this zero the relative radiative corrections blow up.

mind that these factors are a measure for the necessary reparameterization of the GPD when one includes the next order in a given scheme. In Fig. 1 we depict for the typical kinematics in fixed target experiments, i.e., $0.05 \lesssim \xi \lesssim 0.3$, the K factors to NLO as dashed and dotted lines for the $\overline{\text{CS}}$ and $\overline{\text{MS}}$ schemes, respectively. We set $\mu = Q$ and independent of the considered approximation we choose $\alpha_s(\mu_r^2 = 2.5 \text{ GeV}^2) = 0.1\pi$. From the panels (a) and (b) it can be realized that in the $\overline{\text{MS}}$ scheme the radiative corrections to the modulus are up to 20% and 30% for $\alpha = 0.5$ ($\Delta^2 = 0$) and $\alpha = -0.1$ ($\Delta^2 = -0.6 \text{ GeV}^2$), respectively. In the $\overline{\text{CS}}$ scheme these radiative corrections are reduced by 30%. Note that such a reduction has been observed in a quite different processes, namely, for the photon-to-pion transition form factor³. The relative radiative corrections to the phase is in both cases of about 15% at $\xi = 0.6$ and diminishes with decreasing ξ . These findings qualitatively agree with previous ones in which the Radyushkin ansatz for GPDs was used [14].

We study now the radiative corrections to NNLO accuracy. To simplify their evaluation, we take for $c_j^{(2)}$ a fit, given in Ref. [27], rather than the exact expression. For the β_0 proportional term we have checked that within $|c| = |\Re j| \leq 1/2$, see Mellin–Barnes integral (20), the accuracy is on the level of one per mill or better. Outside this region, the deviation can be larger and one might use Fortran routines [28]. For three quark flavors $n_f = 3$, the same scale settings, and the initial value of α_s as specified above for NLO, we plot in Fig. 1 the ratios (22) in the $\overline{\text{CS}}$ scheme for $P = 2$ as solid line. The ξ –dependence of the modulus K –factor, see panels (a) and (b), is rather flat and the modulus decreases on the 5% level. The radiative correction to the phase is again negligible for smaller values of ξ and increases now only to 5% at $\xi = 0.6$. We note that the β_0 induced corrections are about twice times larger than the remaining ones and are opposite in sign, see dash–dotted line. Remarkably, for $\beta_0 = 0$ we find then an opposite behavior of the K factors as in NLO. This arise from a sign alternating series and as for the photon-to-pion transition form factor, this might be considered as a reminiscence on Sudakov double logs [29].

Let us finally address the modification of the scale dependence due to the higher order corrections. Note that the analysis about scheme dependence in Ref. [30] suggests that the discrepancy between the $\overline{\text{CS}}$ and $\overline{\text{MS}}$ schemes at NLO accuracy are mainly induced by the Wilson–coefficients while the evolution yields minor differences. So we only consider here the $\overline{\text{CS}}$ scheme and analogously as in Eq. (22), we quantify the relative changes of $d\mathcal{H}/d\ln Q^2$ by the ratios

$$\dot{K}_{\text{abs}}^P = \left| \frac{d\mathcal{H}^{\text{N}^{\text{PLO}}}}{d\ln Q^2} \right| \bigg/ \left| \frac{d\mathcal{H}^{\text{N}^{\text{P-1LO}}}}{d\ln Q^2} \right|, \quad \dot{K}_{\text{arg}}^P = \left[\pi + \arg \left(\frac{d\mathcal{H}^{\text{N}^{\text{PLO}}}}{d\ln Q^2} \right) \right] \bigg/ \left[\pi + \arg \left(\frac{d\mathcal{H}^{\text{N}^{\text{P-1LO}}}}{d\ln Q^2} \right) \right]. \quad (23)$$

Both the numerator and denominator in the latter $\dot{K}$ –factor are now defined in the interval $[0, 2\pi]$

³This process is also evaluated within the OPE and the reduction of NLO corrections has a common origin. Namely, in the $\overline{\text{CS}}$, compared to the $\overline{\text{MS}}$, scheme the first few Wilson–coefficients are smaller, while the stronger logarithmic growing with increasing j is anyway suppressed by the non–perturbative input, see Eqs. (13) and (19).




Figure 2: The relative change of scale dependence, cf. Eq. (23), in the $\overline{\text{CS}}$ scheme at NLO (dashed, dotted) and NNLO (solid, dash-dotted) versus ξ is depicted for the modulus (a) and phase (b) of the CFF (20) with $\alpha = 0.5$ (dashed, solid) and $\alpha = -0.1$ (dotted, dash-dotted) and $Q^2 = 4 \text{ GeV}^2$. We set $\mu = Q$, $\alpha_s(\mu_r^2 = 2.5 \text{ GeV}^2)/\pi = 0.1$ and took the input (21) at the scale $Q_0^2 = 0.5 \text{ GeV}^2$.

and so the appearance of a zero in the denominator is avoided, cf. footnote 2. We take the same scale setting and initial condition for the (exact) evolution of $\alpha_s(Q)$ as above. The conformal moments (21) are evolved in the $\overline{\text{CS}}$ scheme. The input scale is now $\mu_0^2 = 0.5 \text{ GeV}^2$, which is typical for the matching of perturbative and non-perturbative QCD. The non-leading logs in the solution of the evolution equation (18) are expanded with respect to α_s and are consistently combined with the Wilson-coefficients (12) in the considered order. Here the forward anomalous dimensions are taken from Ref. [31] and the unknown NNLO mixing term $\Delta_{jk}^{\overline{\text{CS}}}$ in Eq. (18) is neglected. This mixing can be suppressed at the input scale within an appropriate initial condition and so we expect only a minor numerical effect; see also Ref. [30]. The dashed and dotted lines in Fig. 2 show that in NLO the scale dependence changes of about 30% to 50% for the modulus and up to 20% for the phase. The latter $\dot{K}$ -factor becomes close to one for smaller values of ξ . The NNLO corrections, compared to the full NLO result, are milder. We observe (solid and dash-dotted line) a 20% or less and a maximal 10% correction for the modulus and phase, respectively.

4 Conclusions

In this letter we have taken the first steps in the full investigation of radiative NNLO corrections to the DVCS scattering amplitude. Thereby, we employed the COPE, which allows an economical treatment of perturbative corrections, and restricted ourselves to the CFF $\mathcal{H}$ in the flavor non-singlet sector. This study can be immediately adopted for $\mathcal{E}$. The extension to the axial vector case, i.e., to $\tilde{\mathcal{H}}$ and $\tilde{\mathcal{E}}$, is straightforward, however, here the anomalous dimensions to three loop order cannot be borrowed from the polarized DIS results. The conformal approach can be also

straightforwardly extended to the flavor singlet sector, which will be presented somewhere else.

We relied on GPDs for which the expansion of the conformal moments in powers of ξ^2 is meaningful and used in our analysis only the leading term. Whether this assumption is too restrictive is an open problem, which might be resolved by lattice calculations [22]. At least there is a warning from perturbative QCD. Namely, in the $\overline{\text{MS}}$ scheme ξ^2 -suppressed contributions in the Wilson-coefficients are resummed and lead to larger radiative corrections than in the $\overline{\text{CS}}$ one. Let us summarize our numerical findings for fixed target kinematics, i.e., $0.05 \lesssim \xi \lesssim 0.3$:

- Compared to the $\overline{\text{MS}}$ scheme, in the $\overline{\text{CS}}$ one the radiative NLO corrections to the modulus of the CFF are reduced by 30% or so, while for the phase no significant differences appear.
- The relative NNLO corrections compared to the NLO ones are of the 5% level or below, where the β_0 proportional ones dominate and determine the sign.
- The change of the scale dependence due to the NLO corrections is rather large and is for the modulus (phase) of the CFF of about 30% to 40% (15% or smaller). Comparing NNLO to NLO, the two-loop corrections are reduced to less than 20% (10%) .

Let us add that the Wilson-coefficients in the vector and axial-vector channel possess a similar j dependence. So we might expect that for $\tilde{\mathcal{H}}$ and $\tilde{\mathcal{E}}$ (axial-vector case) comparable radiative corrections appear. We conclude that our findings support the perturbative description of the DVCS process at scales which are accessible in the kinematics of present fixed target experiments.

Acknowledgement

For discussions I am indebted to A. Manashov, B. Melić, K. Kumerički, K. Passek-Kumerički, and A. Schäfer. This project has been supported by the Deutsche Forschungsgemeinschaft and the U.S. National Science Foundation under grant no. PHY-0456520.

References

- [1] D. Müller, D. Robaschik, B. Geyer, F.-M. Dittes, and J. Hořejši, Fortschr. Phys. **42**, 101 (1994), hep-ph/9812448.
- [2] X. Ji, Phys. Rev. Lett. **78**, 610 (1997), hep-ph/9603249; A. Radyushkin, Phys. Lett. **B380**, 417 (1996), hep-ph/9604317; J. Collins, L. Frankfurt, and M. Strikman, Phys. Rev. **D56**, 2982 (1997), hep-ph/9611433.
- [3] G.P. Lepage and S.J. Brodsky, Phys. Rev. **D22**, 2157 (1980).

- [4] M. Diehl, Phys. Rept. **388**, 41 (2003), hep-ph/0307382; A. V. Belitsky and A. V. Radyushkin, *Unraveling hadron structure with generalized parton distributions*, hep-ph/0504030.
- [5] A. Airapetian *et al.* [HERMES Collaboration], Phys. Rev. Lett. **87** 182001 (2001), hep-ex/0106068; S. Stepanyan *et al.* [CLAS Collaboration], Phys. Rev. Lett. **87** 182002 (2001), hep-ex/0107043; C. Adloff *et al.* [H1 Collaboration], Phys. Lett. **B517**, 47 (2001), hep-ex/0107005; S. Chekanov *et al.* [ZEUS Collaboration], Phys. Lett. **B573** 46 (2003), hep-ex/0305028; C. Aktas *et al.* [H1 Collaboration], Eur. Phys. J. **C44**, 1 (2005), hep-ex/0505061.
- [6] M. Diehl, T. Gousset, B. Pire, and J. P. Ralston, Phys. Lett. **B411**, 193 (1997), hep-ph/9706344.
- [7] A. Belitsky, D. Müller, L. Niedermeier, and A. Schäfer, Nucl. Phys. **B593**, 289 (2001), hep-ph/0004059.
- [8] A. V. Belitsky, D. Müller, and A. Kirchner, Nucl. Phys. **B629**, 323 (2002), hep-ph/0112108.
- [9] D. Müller and A. Kirchner, Eur. Phys. J. **C32**, 347 (2003), hep-ph/0302007, hep-ph/0302007.
- [10] J. Collins and A. Freund, Phys. Rev. **D59**, 074009 (1999), hep-ph/9801262; X. Ji and J. Osborne, Phys. Rev. **D58**, 094018 (1998), hep-ph/9801260; A. Radyushkin, Phys. Rev. **D56**, 5524 (1997), hep-ph/9704207.
- [11] N. Isgur and C.H. Llewellyn Smith, Nucl. Phys. **B317**, 526 (1989).
- [12] A. Belitsky and D. Müller, Phys. Lett. **B417**, 129 (1997), hep-ph/9709379.
- [13] L. Mankiewicz, G. Piller, E. Stein, M. Vanttinen and T. Weigl, Phys. Lett. B **425**, 186 (1998), hep-ph/9712251.
- [14] A. Belitsky, D. Müller, L. Niedermeier, and A. Schäfer, Phys. Lett. **B474**, 163 (2000), hep-ph/9908337.
- [15] A. Freund and M. McDermott, Phys. Rev. **D65**, 074008 (2002), hep-ph/0106319; Eur. Phys. J. **C23**, 651 (2002), hep-ph/0111472.
- [16] D. Müller, Phys. Rev. **D58**, 054005 (1998), hep-ph/9704406.
- [17] V. M. Braun, G. P. Korchemsky, and D. Müller, Prog. in Part. and Nucl. Phys. **51**, 312 (2003), hep-ph/0306057.
- [18] B. Melić, D. Müller, and K. Passek-Kumerički, Phys. Rev. **D68**, 014013 (2003), hep-ph/0212346.

- [19] S. Brodsky, Y. Frishman, G. Lepage, and C. Sachradja, Phys. Lett. **91B**, 239 (1980); A. Efremov and A. Radyushkin, Phys. Lett. **94B**, 245 (1980).
- [20] D. Müller, Phys. Rev. **D49**, 2525 (1994), hep-ph/9411338.
- [21] S. Ferrara, , R. Gatto, and A. Grillo, Phys. Lett. **36B**, 124 (1971), **38B**, 188 (E); ibid. **38B**, 333 (1972); Phys. Rev. **D5**, 3102 (1972); S. Ferrara, A. Grillo, and G. Parisi, Nucl. Phys. **B54**, 552 (1973).
- [22] LHPC, P. Hagler *et al.*, Phys. Rev. **D68**, 034505 (2003), hep-lat/0304018; LHPC and SESAM, P. Hägler *et al.*, ibid. **93**, 112001 (2004), hep-lat/0312014; QCDSF, M. Gockeler *et al.*, Phys. Rev. Lett. **92**, 042002 (2004), hep-ph/0304249.
- [23] D. Müller and A. Schäfer, *Complex conformal spin partial wave expansion of generalized parton distributions and distribution amplitudes*, hep-ph/0509204.
- [24] A. Belitsky and A. Schäfer, Nucl. Phys. **B527**, 235 (1998), hep-ph/9801252.
- [25] G. Curci, W. Furmanski, and R. Petronzio, Nucl. Phys. **B175**, 27 (1980).
- [26] E. B. Zijlstra and W. L. van Neerven, Nucl. Phys. **B383**, 525 (1992).
- [27] W. L. van Neerven and A. Vogt, Nucl. Phys. **B568**, 263 (2000), hep-ph/9907472.
- [28] J. Blümlein, Comput. Phys. Commun. **133**, 76 (2000), hep-ph/0003100; J. Blümlein and S.-O. Moch, Phys. Lett. **B614**, 53 (2005), hep-ph/0503188.
- [29] I. Musatov and A. Radyushkin, Phys. Rev. **D56**, 2713 (1997), hep-ph/9702443.
- [30] D. Müller, Phys. Rev. **D59**, 116003 (1999), hep-ph/9812490.
- [31] S. Moch, J. A. M. Vermaseren, and A. Vogt, Nucl. Phys. **B688**, 101 (2004), hep-ph/0403192.